

**HUBUNGAN ANTARA BERAT BADAN BAYI BARU LAHIR DAN
PARITAS DENGAN RUPTUR PERINEUM PADA PERSALINAN
SPONTAN
DI RSIA BUNDA ARIF PURWOKERTO
TAHUN 2010**

Oleh:

Lysa Destiati¹ dan Fitria Prabandari²,
^{1,2} *Akademi Kebidanan YLPP Purwokerto*
Telp 081326666703, email fitriaoptimisselalu@yahoo.co.id

ABSTRACT

Postpartum haemorrhage can happen by hurt of path delivery, that one of them can from rupture perineum. Rupture can happen at first delivery and not rarely at the next delivery. The causes of rupture perineum's incidence, one of it by a big of newborn baby weight and parities. This research purpose to know a relationship between a newborn baby weight and parities with rupture perineum at spontaneous delivery at RSIA Bunda Arif Purwokerto in 2010. This research was descriptive correlation research with approachment cross sectional. The population of this research was all of spontaneous delivery mother at RSIA Bunda Arif and the sample used was purposive sampling that amount 252 peoples. The method of data analyze used Chi Square test. The known research results: 1) The most baby who born with spontaneous delivery was a enough weight newborn baby (BBLC) as much as (77 %). 2) The most of spontaneous delivery mother was delivery mother who multipara as much as (55,20%). 3) Spontaneous delivery mother who delivered, the most of them experience rupture perineum that was as much as (67,1%). 4) The result of newborn baby weight and rupture perineum's statistic analyze obtained p-value as many as $0,000 < 0,05$ that means H_0 pushed away and H_a acceptance, while the result of parities and rupture perineum's analyze obtained p-value as many as $0,000 < 0,05$ that means H_0 pushed away and H_a accepted. There is a relationship between a newborn baby weight and parities with rupture perineum at spontaneous delivery.

Keywords : *Newborn baby weight, parities, and rupture perineum*